



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2009

---

## **Combination of numerical dating techniques using $^{10}\text{Be}$ in rock boulders and $^{14}\text{C}$ in resilient soil organic matter for reconstructing glacial and periglacial processes in a high alpine catchment during the late Pleistocene and early Holocene**

Favilli, F ; Egli, Markus ; Brandova, D ; Ivy-Ochs, S ; Kubik, P W ; Maisch, M ; Cherubini, P ;  
Haeberli, W

**Abstract:** Glacier fluctuations and paleoclimatic oscillations during the Late Quaternary in Val di Rabbi (Trentino, northern Italy) were reconstructed using a combination of absolute dating techniques ( $^{14}\text{C}$  and  $^{10}\text{Be}$ ) and soil chemical characterization. Extraction and dating of the stable fraction of soil organic matter (SOM) gave valuable information about the minimum age of soil formation and contributed to the deciphering of geomorphic surface dynamics. The comparison of  $^{10}\text{Be}$  surface exposure dating (SED) of rock surfaces with the  $^{14}\text{C}$  ages of resilient (resistant to  $\text{H}_2\text{O}_2$  oxidation) soil organic matter gave a fairly good agreement, but with some questionable aspects. It is concluded that, applied with adequate carefulness, dating of SOM with  $^{14}\text{C}$  might be a useful tool in reconstructing landscape history in high Alpine areas with siliceous parent material. The combination of  $^{14}\text{C}$  dating of SOM with SED with cosmogenic  $^{10}\text{Be}$  (on moraines and erratic boulders) indicated that deglaciation processes in Val di Rabbi were already ongoing by around 14,000 cal BP at an altitude of 2300 m asl and that glacier oscillations might have affected the higher part of the region until about 9000 cal BP.  $^{10}\text{Be}$  and  $^{14}\text{C}$  ages correlate well with the altitude of the sampling sites and with the established Lateglacial chronology.

DOI: <https://doi.org/10.1017/S0033822200055910>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-5045>

Journal Article

Accepted Version

Originally published at:

Favilli, F; Egli, Markus; Brandova, D; Ivy-Ochs, S; Kubik, P W; Maisch, M; Cherubini, P; Haeberli, W (2009). Combination of numerical dating techniques using  $^{10}\text{Be}$  in rock boulders and  $^{14}\text{C}$  in resilient soil organic matter for reconstructing glacial and periglacial processes in a high alpine catchment during the late Pleistocene and early Holocene. *Radiocarbon*, 51(2):537-552.

DOI: <https://doi.org/10.1017/S0033822200055910>

# COMBINATION OF NUMERICAL DATING TECHNIQUES USING $^{10}\text{Be}$ IN ROCK BOULDERS AND $^{14}\text{C}$ IN RESILIENT SOIL ORGANIC MATTER FOR RECONSTRUCTING GLACIAL AND PERIGLACIAL PROCESSES IN A HIGH ALPINE CATCHMENT DURING THE LATE PLEISTOCENE AND EARLY HOLOCENE

Filippo Favilli<sup>1</sup> · Markus Egli<sup>1\*</sup> · Dagmar Brandova<sup>1</sup> · Susan Ivy-Ochs<sup>2</sup> · Peter W. Kubik<sup>3</sup> · Max Maisch<sup>1</sup> · Paolo Cherubini<sup>4</sup> · Wilfried Haeberli<sup>1</sup>

Radiocarbon (Accepted for publication)

**ABSTRACT.** Glacier fluctuations and palaeoclimatic oscillations during the Late Quaternary in Val di Rabbi (Trentino, Northern Italy) were reconstructed using a combination of absolute dating techniques ( $^{14}\text{C}$  and  $^{10}\text{Be}$ ) and soil chemical characterisation. Extraction and dating of the stable fraction of soil organic matter (SOM) could give valuable information about the minimum age of soil formation and contributed to decipher geomorphic surface dynamics. The comparison of surface exposure dating (SED) of rock surfaces with the  $^{14}\text{C}$  ages of resilient (resistant to  $\text{H}_2\text{O}_2$  oxidation) soil organic matter gave a fairly good agreement, but with some questionable aspects. It is concluded that, applied with adequate carefulness, dating of SOM with  $^{14}\text{C}$  might be a useful tool in reconstructing landscape history in high Alpine areas with siliceous parent material. The combination of  $^{14}\text{C}$  dating of SOM with SED of cosmogenic  $^{10}\text{Be}$  (on moraines and erratic boulders) indicated that deglaciation processes in Val di Rabbi were already ongoing around 14000 cal BP at an altitude of 2300 m asl and that glacier oscillations might have affected the higher part of the region until about 9000 cal BP.  $^{10}\text{Be}$  and  $^{14}\text{C}$  ages correlate well with the altitude of the sampling sites and with the established Lateglacial chronology.

## INTRODUCTION

---

<sup>1</sup>Department of Geography, University of Zurich-Irchel, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland.

\*Corresponding author. Email: markus.egli@geo.uzh.ch

<sup>2</sup>Institute of Particle Physics, ETH Zurich, CH-8093 Zurich, Switzerland.

<sup>3</sup>Paul Scherrer Institute, c/o Institute of Particle Physics, ETH Zurich, CH-8093 Zurich, Switzerland.

<sup>4</sup>Swiss Federal Institute for Forest, Snow and Landscape Research, WSL, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland.

Clear evidences of distinct climatic conditions, such as moraines and rockglaciers, make high mountain areas unique "storytellers" about archives of past climate change effects on landscape dynamics. Since the initial work of Penck and Brückner (1901/1909) on Alpine glaciations and the structure of glacier retreat at the end of the last glaciation (the Würmian), numerous authors have worked on ice-age glacial stratigraphy (e.g. Schlüchter 1988, 2004; Keller and Krayss 1987, 2005), on the Lateglacial ice decay (e.g. Maisch 1981; Maisch 1987; Schoeneich 1999; Kerschner et al. 1999; Ivy-Ochs et al. 2004, 2007) and also in detail on Holocene glacier fluctuations in the Alps (e.g. Holzhauser 1984; Hormes et al. 2007). While the general sequence and the related morphostratigraphic positions of the various stadials are quite widely accepted, reliable dating exist only for a few of them (Heitz et al. 1982; Maisch 1987; Schlüchter 1988; Kerschner et al. 1999). This lack of direct in situ ages dating is due to the various limitations of the applied dating methods and to the rareness of optimal sampling sites (representative of the investigated site). The link between distinct moraine sequences and the absolute time scale is still highly speculative, especially in the central Eastern Alps. The time scales of the formation of the landscape reach as far back as the alpine orogeny but mainly relate to the Lateglacial ice retreat at the end of the last Ice Age (20000-11500 yr BP) and include also the entire Holocene period. The Alpine area offers interesting geomorphological settings with pronounced glacial and periglacial features such as Lateglacial moraines, erratic boulders and rock glaciers that give the opportunity to date the retreat stages of the glaciers and to investigate the subsequent geomorphological modifications during the Holocene. Surface exposure dating (SED) is based on the in-situ production and accumulation of so-called cosmogenic nuclides (e.g.  $^{10}\text{Be}$ ,  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ ) within the first few decimetres of an exposed rock surface (Lal 1991; Gosse and Phillips 2001). This accumulation is used to determine the elapsed time since the rock surface was exposed to cosmic rays (Nishiizumi et al. 1989; Gosse and Phillips 2001). Hence, the deposition age of boulders in various topographical situations can be measured. Several studies have shown that SED is an innovative and powerful tool for the reconstruction of glacial and climate history (e.g. Ivy-Ochs et al. 2004; Briner et al. 2005; Owen et al. 2005; Shulmeister et al. 2005; Smith et al. 2005). In order to compare the magnitude and timing of former glacier extents, absolute dating methods can be applied and compared to determine the age of particular glacial stages and – in combination with other approaches, i.e. by palaeoglaciological reconstructions (Equilibrium Line Altitude (ELA) depressions) – the palaeoclimatic signal they represent (Kelly et al. 2004).

Soils can provide useful information for the geomorphological interpretation of mountainous terrain. Their spatial distribution in Alpine areas reflects the impact of the soil-forming factors and, once their influence is known, soils can help to introduce additive aspects of landscape evolution

that otherwise may be left undetected (Birkeland et al. 2003). The dating of soil organic matter (SOM) with  $^{14}\text{C}$  is well suitable, if humified and stable substances which were produced almost at the beginning of soil formation can be found and dated, giving the possibility to elucidate soil dynamic processes (Scharpenseel and Becker-Heidmann 1992).

The main aim of the study was to test and compare signals of landscape evolution obtained from soils developed on moraines using  $^{14}\text{C}$  ages from stable organic matter with SED obtained from rock boulders. For this purpose we investigated soils developed on morainic and glacial substrata (paragneiss) with respect to their chemical composition and to the radiocarbon ages of the most stable resilient organic material (e.g. Mikutta et al. 2006; Eusterhues et al. 2005). In the vicinity of the studied soils we sampled erratic boulders ( $> 2\text{m}^3$ , in stable position) located on lateral and recessional moraines for  $^{10}\text{Be}$  surface exposure dating. The comparison between the  $^{14}\text{C}$  and  $^{10}\text{Be}$  method enabled to test the validity of the dating technique using resilient SOM. The combination of ages derived from  $^{10}\text{Be}$  and  $^{14}\text{C}$  dating should, furthermore, give insights into the deglaciation processes and the climatic fluctuations occurring during the late Pleistocene and early Holocene in the Central-Eastern Alps in Italy.

## SITE DESCRIPTION

The investigation area is located in Val di Rabbi, in a small side-cirque within a lateral valley of Val di Sole, Trentino, in the southern Alpine belt of northern Italy (Figure 1). The climate of the valley ranges from temperate to alpine (above the timberline). Mean annual temperatures range from  $8.2^\circ\text{C}$  (valley floor – at around 800 m asl) to around  $0^\circ\text{C}$  (at 2400 m asl) and mean annual precipitation approximately from 800 to 1300 mm/year (Servizio Idrografico 1959). The timberline is close to 2100-2200 m asl and the forests are dominated by the conifers *Larix decidua* and *Picea abies* (Pedrotti et al. 1974). Areas above 2300 m are covered with rocks, boulders and short-grass meadows dominated by *Carex curvula* and *Nardus stricta*.

The four investigated soils and boulders (Figure 1; Table 1 and Table 2) are located between 2100 m asl and 2456 m asl, hence, close to the timberline and in the high alpine zone (Figure 1; Table 1 and Table 2). The soil types were *Entic Podzol* and *Haplic Podzol* at lower altitudes (between 2000 and 2200 m asl) and *Cambic Leptosol* and *Brunic Regosol* developed at around 2300 m asl according to the WRB (IUSS working group 2006). According to the Soil Taxonomy (Soil Survey Staff 2006), the soil moisture regime is udic (humid conditions,  $<90$  days/year with a dry soil) at all sites and the soil temperature regime is cryic (mean annual temperature  $< 8^\circ\text{C}$ , without permafrost). Soil material was collected from excavated pits and undisturbed soil samples were taken, where

possible, down to the BC or C horizon (parent material). Two to four kilograms of soil material were collected per soil horizon from the four soil pits (Hitz et al. 2002). The landscape near the investigation sites was strongly influenced by former glaciers and the sampled soils developed on moraines or on rock glaciers (paragneiss) being inactive today. According to the geomorphological studies of Baroni and Carton (1990) and Filippi et al. (2007), surface ages were estimated to 14000 to 16000 years. Large lateral moraines extend down to 1700 - 2000 m asl and glacial deposits, such as erratic boulders, can be found up to 2600 m asl. Big boulders ( $> 2\text{m}^3$ ) with prominent quartz veins in flat and stable positions are widespread in the whole investigated area.

## **METHODS**

### **Soil chemistry and physics**

The soil samples were air-dried, large aggregates were gently broken by hand and sieved to  $<2$  mm. Total C and N contents of the soil were measured with a C/H/N analyser (Elementar Vario EL, Elementar Analysensysteme GmbH) using oven-dried ( $105^\circ\text{C}$ ) and ball-milled fine earth. Total C corresponds in our case to organic C due to the absence of any carbonates in the soil. Soil pH (in 0.01 M  $\text{CaCl}_2$ ) was determined on air-dried samples of the fine earth fraction using a soil solution ratio of 1:2.5. After a pre-treatment of the samples with  $\text{H}_2\text{O}_2$  (3%), particle size distribution of the soils was measured by a combined method consisting of sieving the coarser particles (2000 - 32  $\mu\text{m}$ ) and the measurement of the finer particles ( $< 32$   $\mu\text{m}$ ) by means of an X-ray sedimentometer (SediGraph 5100). The dithionite- ( $\text{Fe}_\text{d}$ ,  $\text{Al}_\text{d}$ ) and oxalate-extractable ( $\text{Fe}_\text{o}$ ,  $\text{Al}_\text{o}$ ) iron and aluminium fractions were extracted according to McKeague et al. (1971) and analysed by AAS (Atomic Absorption Spectrometry – AAnalyst 700, Perkin Elmer, USA)

### **Extraction of the stable organic matter**

Acting on the assumption that chemical oxidation mimics natural oxidative processes, we treated the soils with 10%  $\text{H}_2\text{O}_2$  to eliminate the more labile organic material from the more refractory organic matter (Plante et al. 2004; Eusterhues et al. 2005; Mikutta et al. 2006; Helfrich et al. 2007). The stable fraction that remained at the end of the treatment was supposed to be part of the first organic matter formed in the sediment after glacier retreat (Favilli et al. 2008) and to provide minimum ages of deposition of the moraines and of deglaciation. One gram of air-dried soil was wetted for 10 min with few ml of distilled water in a 150-ml beaker. Afterwards, 90 ml of 10%  $\text{H}_2\text{O}_2$  were added. The procedure was run at a minimum temperature of  $50^\circ\text{C}$  throughout the treatment period. The beakers were closed with two layers of parafilm (Pechiney Parafilm M

PM999, Chicago, USA) to avoid evaporation of the reagent. Peroxide treatments were performed for 168 hours (7 days). At the end of the treatment the samples were washed three times with 40 ml deionised water, freeze-dried, weighted, analysed for total C and N and  $^{14}\text{C}$  dated.

### **Radiocarbon dating**

The samples were heated and catalytically reduced over cobalt powder at  $550^{\circ}\text{C}$  to elemental carbon (graphite). This mixture was pressed into a target and the ratios  $^{14}\text{C}:^{12}\text{C}$  (for radiocarbon age) were measured by Accelerator Mass Spectrometry (AMS) using the tandem accelerator of the Institute of Particle Physics at the Swiss Federal Institute of Technology Zürich (ETH). The calendar ages were obtained using the CalPal and OxCal calibration programs (Danzeglocke et al. 2008; Bronk Ramsey 2001).

### **Exposure dating**

Four large boulders with volumes  $> 2\text{m}^3$  were sampled for exposure dating on top of moraines. The assumption is that certain boulders were transported by the glacier during the last glaciation and were, since the time of the last glacier retreat, exposed to cosmic rays without being moved from their present position. Gosse et al. (1995) demonstrated that on a single alpine moraine the cosmogenic nuclide ages reflect the timing of deposition. These boulders were chosen for dating in order to exclude any long-term effects from slope-movement processes (Table 2). The samples were crushed, sieved and leached in order to obtain pure quartz following Kohl and Nishiizumi (1992) and Ivy-Ochs (1996).  $^9\text{Be}$  solution was added to the dried quartz which was then dissolved in 40% HF. The Be was isolated using anion and cation exchange columns followed by selective pH precipitation techniques (Ivy-Ochs 1996). The  $^{10}\text{Be}/^9\text{Be}$  ratios were measured at the ETH AMS facility. The surface exposure ages listed in Table 2 were calculated using a sea-level high-latitude production rate of  $5.1 \pm 0.3$  ( $^{10}\text{Be}$  atoms / g  $\text{SiO}_2$  / year) with a 2.2 % production due to negative muon capture (Stone 2000). This production rate was scaled for latitude (geographic) and altitude (Stone 2000) and corrected for sample thickness assuming an exponential depth profile, a rock density of  $2.65 \text{ g cm}^{-2}$  and an effective collimated radiation attenuation length of  $155 \text{ g cm}^{-2}$  (Gosse and Phillips 2001). Topographic shielding was based on a zenith angle dependence of  $(\sin\theta)^{2.3}$  (Dunne et al. 1999). The errors given for each boulder age reflect the analytical uncertainties of the AMS measurement parameters. The exposure age was corrected for mean snow cover (Gosse and Phillips 2001) and an assumed snow density of  $0.3 \text{ g cm}^{-3}$ . The duration of the snow cover was estimated according to Auer et al. (2003) from climatic data supplied by the Provincia Autonoma di Trento (Dipartimento Protezione Civile e Tutela del Territorio – Ufficio Previsioni e

Organizzazione). The snow correction increases the exposure age by around 10%. A geomagnetic field correction (Pigati and Lifton 2004) was omitted because its possible effect is small (1-2 %).

## RESULTS

### **Chemical composition and physical characteristics of the soils**

The investigated soils were characterised by a high proportion of rock fragments ranging from 0% up to 60% of the total weight (Table 3). These are typical values for alpine soils on a moraine type substratum (Egli et al. 2001). All investigated soils had a sandy to silty-sandy texture. The proportion of sand decreased towards the soil surface, and correspondingly silt and clay increased (Table 3). The decrease of the grain sizes from the bottom to the top of the soil profile is a clear effect of weathering. Due to the siliceous parent material, the soils show pronounced acidification (Table 4). Eluviation and illuviation of Fe and Al was typical for the podzolised soils (S1 and S2). A distinct increase of dithionate and oxalate iron and aluminium was observed in all spodic horizons (Table 4). The soil S3 was rather shallow but showed the typical characteristics of a developing-podzol (Al, Fe and organic C translocation within the soil profile and formation of a Bhs horizon). In the soil profile S4 no clear translocation of Fe and Al forms is detectable due to a lower stage of soil development. This soil developed in an area with former solifluidal activity. The soil had a polygenetic horizon sequence due to the accumulation of younger material on top of a former surface (Figure 2). The concentration of organic C in the profile showed a strong increase in the Ab horizon, thus confirming the macromorphological observation (Table 4).

### **Radiocarbon age of soil organic matter and $^{10}\text{Be}$ age of boulders**

The calibration of the  $^{14}\text{C}$  ages using two different calibration programs (CalPal and OxCal) gave comparable results in all the samples (Table 5). The ages of SOM varied considerably within all the soil profiles (Table 5). The highest age of the  $\text{H}_2\text{O}_2$  treated samples was usually measured in the E or B horizon (Bs or Bhs) and gives an indication about the start of soil formation.  $\text{H}_2\text{O}_2$  is a strong reagent to oxidise all labile organic matter in soils (Favilli et al. 2008). The measured lower ages in the AE horizon of profile S2 and the AE1 horizon of profile S3, compared to those of the subsoil, are most probably due to the very high C content and therefore, either to the incapability of  $\text{H}_2\text{O}_2$  to oxidise all young OM or to young OM which was able to expel older C. The untreated soil organic matter from the whole soil profile S3 (Table 5, Bhs horizon) has in general a very young age. Strong leaching conditions, due to the very low pH, have caused the downward migration of young soluble organic material compounds with soil percolating water into the subsoil. The  $\text{H}_2\text{O}_2$ -

treatment, performed on the sample from the Bhs horizon gave an age of  $10690 \pm 110$  cal BP of the resilient OM-fraction.

It seems that at least a part of the oldest fraction is still present in the top- and subsoil and that the treatment with hydrogen peroxide is able to detect some of the initially formed OM. The  $^{14}\text{C}$  results show that the site S4 has a polygenetic soil. Soil formation in the initial layer started around 13800 cal BP and ended, due to accumulation of eroded material, probably slope deposits, around 2600 cal BP (Table 5). This burial event was inferred from dating root remains in the Ab horizon. The accumulation was probably due to periglacial activity/processes (solifluction) during a colder period around 3000 yr BP. This periglacial period would fit with the 'Göschener' cold phase I, which occurred around 3000 cal BP (Maisch et al. 1999; Figure 4). Preweathered sediments containing already organic material (having a maximum age of 8700-8900 cal BP) were deposited on top of the original soil (horizon A/Bw2; Figure 2).

The  $^{10}\text{Be}$  dates from the sampled boulders range from 13230 yr to 8840 yr (Table 2). The  $^{10}\text{Be}$  ages decrease with increasing altitude which is in agreement with the general direction of retreat of the glaciers after the LGM (Last Glacial Maximum). The position of the boulders in the investigation area supports the calculated ages (Figure 1 and Figure 5). If exposure ages and  $^{14}\text{C}$  ages (start of soil formation) are plotted versus elevation, a significant ( $r^2 = 0.68$ ;  $p = 0.0115$ ), linear relationship is obtained (Figure 3).

According to the radiocarbon age of stable organic matter extracted from soil S1, the oldest moraine (at an altitude of 2100 m asl) was already deposited around 17000 cal BP (Figure 4 and Table 5). The soil S4 at 2370 m asl started to develop 13840 cal BP or earlier. More or less simultaneously the boulder B1, located at 2247 m asl, was deposited on a moraine. Around 11700 yr ago, the site B2 at 2360 m asl was affected by the readvance of a small glacier tongue, most probably during the early Preboreal (Ivy-Ochs et al. 2006a) or probably late Egesen glacial stage (corresponding to Younger Dryas) with the deposition of new material (Figure 5). The soil S3, located in short distance from B2, started its evolution shortly after the deposition of the moraine. The chemical evolution of the soil profile S3 confirms that this soil has had an undisturbed development during the last 10000 years – a time span which leads to well-developed podzols (e.g. Starr 1991; Barrett and Schaetzl 1992; Lundström et al. 2000). The following period, the Holocene, was characterised by higher temperatures (Clapperton 1995) which finally caused the total disappearance of the glaciers in the investigation area. Periglacial activity (solifluction) was, however, traceable at the site S4 until around 2600 cal BP. This soil profile is polygenetic and strongly influenced by solifluction (accumulation of soil material because of slope deposits, burial of a former soil) around that period.



## DISCUSSION

The moraines in the study area are characterised by sharp crests and numerous large boulders (see also Hormes et al. 2007 for moraine description). The investigated soils are well developed and show the typical characteristics of podzolisation, with the formation of an eluvial and a illuvial horizon. The oxidation of SOM with  $\text{H}_2\text{O}_2$  left behind intrinsically resistant soil organic matter (Theng et al. 1992; Cuypers et al. 2002; Favilli et al. 2008). Humic compounds may form close and strong associations with the mineral phase, especially with clay minerals (Righi and Meunier 1995). Clay organic complexes, once formed, do not easily exchange the organic component with infiltrated younger humus components (Scharpenseel and Becker- Heidmann 1992). A higher  $^{14}\text{C}$  age in the topsoil or near the surface should be the theoretical expected since soil formation starts at the surface and proceeds with time into greater depths. Organic matter from plants and animals is incorporated into soil material and mixed with the mineral part components. Because the investigated soils are podzolised and consequently have a low biological activity and no earthworms, the highest ages of the stable SOM should be found in the topsoil. Such a typical trend was found in profile S1. In contrast to the treated samples, the untreated samples (S1) showed an increase of age with soil depth. This was explained by a continuous rejuvenation of SOM from the soil surface. Most of modern carbon is less than 100 years old and decreases exponentially with increasing depth leading to an increase of the percentage of old carbon present (O'Brien and Stout 1978; Mikutta et al. 2006). On one hand, it seems that at least a part of the oldest fraction is still present in the top- and subsoil and that the treatment with hydrogen peroxide is able to detect some of the initially formed OM. On the other hand, it does not seem that in every case the oldest organic material can be found in the topsoil. This means that the top- and subsoil material has to be dated to obtain the oldest SOM fraction. Radiocarbon ages prove the late Pleistocene and Holocene age of the surfaces. The  $^{14}\text{C}$  ages of resilient SOM fit well to the  $^{10}\text{Be}$  ages of rock boulders. Radiocarbon dating of stable organic matter therefore seems to be a promising tool for dating Holocene and late Pleistocene-aged soils. The significant linear correlation found between the  $^{14}\text{C}$  and  $^{10}\text{Be}$  ages and the altitude of the investigated sites enables hypotheses about the chronology of glacial and periglacial processes that occurred in the investigated area. The combination of these two dating techniques, therefore, helps to decipher landscape history of high Alpine regions on siliceous parent material in the late Pleistocene/Holocene.

The investigation area experienced deglaciation between 18000 and 9000 cal BP. The ages of the soils and of the exposed boulders reflect quite clearly the movement and the timing of the glacier retreat and – in a general sense – the dynamics of Alpine landscape geomorphology. The ages of the

soils and of the exposed rocks correspond quite well with the general chronology of the Lateglacial and the Holocene period, even if some aspects remain open. Deglaciation from the LGM occurred worldwide during the three millennia 18 – 15 ka BP, documented by ice cores and global sea level records (e.g. Fairbanks 1989; Johnsen et al. 1992). Studies from the Upper Engadine (Switzerland) confirm that the retreat of the main valley glacier was substantially progressed already at 16000 cal BP (Studer 2005). This fits well with the radiocarbon age of stable organic matter extracted from the soil S1 having an age of around 17000 cal BP.

The overall trend of deglaciation continued after ca. 15 ka BP but was interrupted by extensive glacier readvance between 15.4 and 13.2 <sup>14</sup>C uncal. ka BP (Gschnitz stadial; Kohl 1986; Ivy-Ochs et al. 2006b) and reflects an ELA 650-700m below that of the Little Ice Age reference ELA (Ivy-Ochs et al. 2006b). The age of 14683 ± 313 cal BP in the topsoil of profile S1 could point to a readvance of the glaciers and deposition of moraine material in the Older or Oldest Dryas. In this case, profile S1 would be polygenetic (burial of a pre-existing soil surface with the start of a new soil development). A polygenetic soil is, however, rather improbable as no signs of accumulation and burial were visible and measurable e.g. by a distinct increase of rock fragments (material > 2mm) near the surface (see Table 3). Until the end of the Gschnitz stadial the oscillations of the glaciers occurred down to an altitude of around 1700 m asl. The period between about 15.5 and 12.8 ka cal BP has been recognized as the Bølling-Allerød interstadial (Alley et al. 1993; Maisch et al. 1999; Schaub et al. 2008). <sup>14</sup>C dates with varve counting indicated that the Upper Engadine lakes became ice-free prior to approximately 13500 <sup>14</sup>C uncal BP (around 16000 cal BP) (Ohlendorf 1998). During this period of warming, Val di Rabbi also experienced a further retreat of its glaciers in a relatively short period of time compared to the overall deglaciation.

After the “Younger Dryas” (about 13.0 – 11.6 ka cal BP; e.g. Lundqvist 1986; Maisch et al. 1999; Studer 2005) soils started to develop also at higher altitudes (e.g. profile S2).

The boulders sampled on the most elevated moraine have an age between 8840 ± 690 - 9920 ± 780 yr. Their average age points to around 9500 yr. They were probably deposited during a readvance phase in the Boreal (Oberhalbstein/Schams cold phase?). Periglacial activity measured probably increased around 2594 ± 101 cal BP (profile S4). This would fit with the ‘Göschener’ cold phase I which occurred around 3000 cal BP (Figure 4). Preweathered sediments containing already organic material (having a maximum age of 8700-8900 cal BP) were deposited on top of the original soil (Figure 2). According to previous studies (Egli et al. 2001) the formation and development of podzolisation features at this altitude takes at least 1000-3000 years. Typical podzolisation features did not develop in these 2600-year-old slope deposits. 2600 years of soil evolution were obviously not a sufficiently long time for the development of podzolic characteristics.

## CONCLUSIONS

We compared and combined radiocarbon ages of stable organic matter in four soil profiles with the  $^{10}\text{Be}$  exposure ages of erratic boulders in order to reconstruct the timing of deglaciation at the transition of the Pleistocene/Holocene. We obtained the following principle findings:

- The  $^{14}\text{C}$  data of the resilient organic matter in soils fit quite well to and can be combined with the  $^{10}\text{Be}$  age sequence obtained from rock boulders.
- Extraction and dating of stable organic matter from alpine soils gives a numerical value about an ice-free minimum age for deglaciation of land surfaces and the beginning of soil formation. More research is, however, needed to confirm these results.
- Dating of resilient OM in soils offers new perspectives in deciphering landscape history but has to be critically evaluated and discussed.
- $^{10}\text{Be}$  and  $^{14}\text{C}$  ages correlated with elevation of the investigated site and with the chronology of the Lateglacial /early Holocene re-advance and retreat phases of glaciers.
- The combination of  $^{10}\text{Be}$  and  $^{14}\text{C}$  dating techniques, applied in relatively small catchments in Alpine environments, despite the limited number of samples, seems to be a promising tool for a better understanding of the geomorphology and palaeoclimate.
- Obtained ages reveal the chronology of the glacial and periglacial processes that occurred in Val di Rabbi between 18000 and 9000 years BP.

## ACKNOWLEDGEMENTS

The research was supported by a grant from the “Stiftung für wissenschaftliche Forschung” of the University of Zurich. We are indebted to I. Woodhatch and B. Kägi for the support in the laboratory. We also would like to thank Dr. Sauer and Dr. Szwarczewski for their helpful comments on an earlier version of the manuscript.

## REFERENCES

- Alley RB, Meese DA, Shuman CA. 1993. Abrupt increase in Greenland snow accumulation at the end of the Younger Dryas event. *Nature* 362: 527-529.
- Auer I, Böhm R, Potzmann P, Ungersböck M. 2003. Änderung der Frosthäufigkeit in Österreich. *Extended Abstract of 6<sup>th</sup> Deutsche Klimatagung*, 22-25 Sept. 2003, Postdam, BRD.

- Barett LR, Schaetzl RJ. 1992. An examination of podzolization near Lake Michigan using chronofunctions. *Canadian Journal of Soil Science* 72: 527-541.
- Baroni C, Carton A. 1990. Variazioni oloceniche della Vedretta della Lobbia (gruppo dell'Adamello, Alpi Centrali). *Geografia Fisica e Dinamica Quaternaria* 13: 105-119.
- Birkeland PW, Shroba RR, Burns SF, Price AB, Tonkin PJ. 2003. Integrating soils and geomorphology in mountains – an example from the Front Range of Colorado. *Geomorphology* 55: 329-344.
- Briner JP, Kaufman DS, Manley WF, Finkel RC, Caffee MW. 2005. Cosmogenic exposure dating of late Pleistocene moraine stabilization in Alaska. *GSA Bulletin* 117: 1108–1120.
- Bronk Ramsey, C, 2001, Development of the radiocarbon calibration program OxCal, *Radiocarbon*, 43: 355-363
- Clapperton C. 1995. Fluctuations of local glaciers at the termination of the Pleistocene: 18-8 ka BP. *Quaternary International* 28: 41-50.
- Cuypers C, Grotenhuis T, Nierop KGJ, Franco EM, de Jager A, Rulkens W. 2002. Amorphous and condensed organic matter domains: the effect of persulfate oxidation on the composition of soil/sediment organic matter. *Chemosphere* 48: 919-931.
- Danzeglocke U, Jöris O, Weninger B. 2008. CalPal-2007online. <http://www.calpal-online.de/>, accessed 2008-03-12.
- Dunne J, Elmore D, Muzikar P. 1999. Scaling factors for the rates of production of cosmogenic nuclides for geometric shielding and attenuation at depth on sloped surfaces. *Geomorphology* 27: 3-11
- Egli M, Mirabella A, Fitze P. 2001. Weathering and evolution of soils formed on granitic, glacial deposits: results from chronosequences of Swiss Alpine environments. *Catena* 45: 19-47.
- Eusterhues K, Rumpel C, Kögel-Knabner I. 2005. Stabilization of soil organic matter isolated via oxidative degradation. *Organic Geochemistry* 36: 1567-1575.
- Fairbanks RG. 1989. A 17,000-year glacio-eustatic sea level record: influence of glacial melting rates on the Younger Dryas event and deep ocean circulation. *Nature* 347: 637-642.
- Favilli F, Egli M, Cherubini P, Sartori G, Delbos E, Haeberli W. 2008. Comparison of different methods of obtaining a resilient organic matter fraction in Alpine soils. *Geoderma* 145: 355-369.
- Filippi ML, Arpent E, Heiri O, Frisia S, Angeli N, van der Borg K, Blockley S. 2007. Lake Lavarone Late-glacial to present palaeoenvironmental changes: a unique multi-proxy record from Trentino, NE Italy. *Geophysical Research Abstracts* 9: 06639.
- Gosse JC, Klein J, Evenson EB, Lawn B, Middleton R. 1995. Beryllium-10 dating of the duration and retreat of the last Pinedale glacial sequence. *Science* 268: 1329-1333.

- Gosse JC, Phillips FM. 2001. Terrestrial in situ cosmogenic nuclides: theory and application, *Quaternary Science Review* 20: 1475–1560.
- Heitz A, Panchakunnel P, Zoller H. 1982. Zum Problem der  $^{14}\text{C}$ -Datierung im Veltlin und Oberengadin. *Physische Geographie* 1: 91-101.
- Helfrich M, Flessa H, Mikutta R, Dreves A, Ludwig B. 2007. Comparison of chemical fractionations methods for isolating stable soil organic carbon pools. *European Journal of Soil Science* 58: 1316-1329.
- Hitz C, Egli M, Fitze P. 2002. Determination of the sampling volume for representative analysis of alpine soils. *Zeitschrift für Pflanzenernährung und Bodenkunde* 165: 326-331.
- Holzhauser H. 1984. Zur Geschichte der Aletschgletscher und des Fieschergletschers. Diss. Univ. Zürich.
- Hormes A, Ivy-Ochs S, Kubik PW, Ferreli L, Michetti AM. 2007.  $^{10}\text{Be}$  exposure ages of a rock avalanche and a late glacial moraine in Alta Valtellina, Italian Alps. *Quaternary International*, In Press. doi: 10.1016/j.quaint.2007.06.036.
- IUSS Working Group WRB. 2006. World Reference Base for Soil Resources 2006, 2<sup>nd</sup> edition, World Soil Resources Reports No. 103, FAO (Food and Agriculture Organisation of the United Nations), Rome.
- Ivy-Ochs S. 1996. The dating of rock surfaces using in situ produced  $^{10}\text{Be}$ ,  $^{26}\text{Al}$  and  $^{36}\text{Cl}$ , with examples from Antarctica and the Swiss Alps. Ph.D. Thesis, ETH Zürich, No. 11763.
- Ivy-Ochs SD, Schäfer J, Kubik PW, Synal HA, Schlüchter C. 2004. Timing of deglaciation on the northern Alpine foreland (Switzerland). *Eclogae Geologicae Helveticae* 97: 47-55.
- Ivy-Ochs SD, Kerschner H, Reuther A, Maisch M, Sailer R, Schaefer J, Kubik PW, Synal HA, Schlüchter C. 2006a. The timing of glacier advances in the northern European Alps based on surface exposure dating with cosmogenic  $^{10}\text{Be}$ ,  $^{26}\text{Al}$ ,  $^{36}\text{Cl}$ , and  $^{21}\text{Ne}$ . In: Siame L.L., Bourlès, D.L., Brown, E.T, editors. *In Situ-Produced Cosmogenic Nuclides and Quantification of Geological Processes*: Geological Society of America Special Paper 415: 43–60.
- Ivy-Ochs SD, Kerschner H, Kubik PW, Schlüchter C. 2006b. Glacier response in the European Alps to Heinrich Event 1 cooling: the Gschnitz stadial. *Journal of Quaternary Science* 21: 115-130.
- Ivy-Ochs SD, Kerschner H, Schlüchter C. 2007. Cosmogenic nuclides and the dating of Lateglacial and Early Holocene glacier variations: The Alpine perspective. *Quaternary International* 164-165: 53-63.
- Johnsen SJ, Clausen HB, Dansgaard W, Fuhrer K, Gundestrup N, Hammer CU, Iversen P, Jouzel J, Stauffer B, Steffensen JP. 1992. Irregular glacial interstadials recorded in a new Greenland ice core. *Nature* 359: 311-313.

- Keller O, Krayss E. 1987. Die hochwürmeiszeitlichen Rückzugsphasen des Rhein-Vorlandgletschers und der erste alpine Eisrandkomplex im Spätglazial. *Geographica Helvetica* 42: 169-178.
- Keller O, Krayss E. 2005. Der Rhein-Linth-Gletscher im letzten Hochglazial: *Vierteljahresschrift der Naturforschenden Gesellschaft in Zürich* (150):19-32 and 69-85.
- Kelly MA, Kubik PW, Von Blanckenburg F, Schlüchter C. 2004. Surface exposure dating of the Great Aletsch Glacier Egesen moraine system, western Swiss Alps, using the cosmogenic nuclide  $^{10}\text{Be}$ . *Journal of Quaternary Science* 19: 431-441.
- Kerschner H, Ivy-Ochs S, Schlüchter C. 1999. Paleoclimatic interpretation of the early late-glacial glacier in the Gschnitz valley, central Alps, Austria. *Annals of Glaciology* 28: 135-140.
- Kohl H. 1986. Pleistocene glaciations in Austria. *Quaternary Science Reviews* 5: 421-428.
- Kohl CP, Nishiizumi K. 1992. Chemical isolation of quartz for measurement of in-situ produced cosmogenic nuclides. *Geochemica et Cosmochimica Acta* 56: 3583-3587.
- Lal D. 1991. Cosmic rays labeling of erosion surfaces: in-situ nuclide production rates and erosion models. *Earth and Planetary Science Letters* 104: 424-439.
- Lundqvist J. 1986. Stratigraphy of the central area of the Scandinavian glaciation. *Quaternary Science Reviews* 5: 251-268.
- Lundström US, van Breemen V, Bain D. 2000. The podzolization process. A review. *Geoderma* 94: 91-107.
- Maisch M. 1981. Glazialmorphologische und gletschergeschichtliche Untersuchungen im Gebiet zwischen Landwasser- und Albulatal (Kt. Graubünden, Schweiz). [dissertation]. University of Zürich (CH): Physische Geographie, vol. 3.
- Maisch M. 1987. Zur Gletschergeschichte des alpinen Spätglazials: Analyse und Interpretation von Schneegrenzdaten. *Geographica Helvetica* 42: 63-71.
- Maisch M, Wipf A, Denzler B, Battaglia J, Benz C. 1999. Die Gletscher der Schweizer Alpen. Gletscherhochstand 1850, aktuelle Vergletscherung, Gletscherschwund-Szenarien. Schlussbericht NFP 31 Projekt, vdf-Hochschulverlag ETH Zürich.
- McKeague JA, Brydon JE, Miles NM. 1971. Differentiation of forms of extractable iron and aluminium in soils. *Soil Science Society of America Proceedings* 35: 33-38.
- Mikutta R, Kleber M, Torn MS, Jahn R. 2006. Stabilization of organic matter: association with minerals or chemical recalcitrance? *Biogeochemistry* 77: 25-56.
- Nishiizumi K, Winterer EL, Kohl CP, Klein J, Middleton R, Lal D, Arnold JR. 1989. Cosmogenic ray production rates of  $^{10}\text{Be}$  and  $^{26}\text{Al}$  in quartz from glacially polished rocks. *Journal of Geophysical Research* 94: 907-915.

- O'Brien BJ, Stout JD. 1978. Movement and turnover of soil organic matter as indicated by carbon isotope measurements. *Soil Biology and Biochemistry* 10: 309-317.
- Ohlendorf C. 1998. High Alpine lake sediments as chronicles for regional glacier and climate history in the Upper Engadine, southeastern Switzerland. [dissertation] No. 12705, ETH Zürich.
- Owen LA, Finkel RC, Barnard PL, Haizhou M, Asahi K, Caffee MW, Derbyshire E. 2005. Climatic and topographic controls on the style and timing of Late Quaternary glaciation throughout Tibet and the Himalaya defined by  $^{10}\text{Be}$  cosmogenic radionuclide surface exposure dating. *Quaternary Science Reviews* 24: 1391–1411.
- Penck A, Brückner E. 1901/09. Die Alpen im Eiszeitalter, Vol. 1-3, Tauchnitz.
- Pedrotti F, Orsomando E, Francalancia C, Cortini Pedrotti C. 1974. Carta della vegetazione del Parco Nazionale dello Stelvio, scala 1:50.000. *Dip. Di Botanica, Università di Camerino (Italia)*.
- Plante AF, Chenu C, Balabane M, Mariotti A, Righi D. 2004. Peroxide oxidation of clay-associated organic matter in a cultivation chronosequence. *European Journal of Soil Science* 55: 471-478.
- Pigati, J.S., Lifton, N.A. 2004. Geomagnetic effects on time-integrated cosmogenic nuclide production with emphasis on in situ C-14 and Be-10, *Earth and Planetary Science Letters* 226: 193-205.
- Righi D, Meunier A. 1995. Origin of clays by rock weathering and soil formation. In: Velde, B, editor. *Origin and Mineralogy of Clays*, Springer-Verlag, Berlin: 43-161.
- Scharpenseel HW, Becker-Heidmann P. 1992. Twenty-five years of radiocarbon dating soils: paradigm of erring and learning. *Radiocarbon* 34: 541-549.
- Schaub M, Büntgen U, Kaiser KF, Kromer B, Talamo S, Andersen KK, Rasmussen SO. 2008. Lateglacial environmental variability from Swiss tree rings. *Quaternary Science Reviews* 27: 29-41.
- Schlüchter C. 1988. The deglaciation of the Swiss Alps: A paleoclimatic event with chronological problems. *Bulletin de l'Association Francaise pour l'étude du Quaternaire* (2/3): 141-145.
- Schlüchter C. 2004. The Swiss glacial record a schematic summary. *Quaternary Glaciations - Extent and Chronology Part I: Europe*. Ehlers J, Gibbard PL, editors. Elsevier: London: 413-418.
- Schoeneich P. 1999. Le retrait glaciaire dans les vallées des Ormonts, de l'Hongrin et de l'Étivaz (Préalpes vaudoises). *Thèse de la Faculté des lettres de l'Université de Lausanne, Travaux et recherche* 14 (1/2).
- Servizio Idrografico. 1959. Precipitazione medie mensili ed annue per il Trentino 1921-1950. *Istituto Poligrafico dello Stato*, Roma.

- Shulmeister J, Fink D, Augustinus PC. 2005. A cosmogenic nuclide chronology of the last glacial transition in North-West Nelson, New Zealand – new insights in Southern Hemisphere climate forcing during the last deglaciation. *Earth and Planetary Science Letters* 233: 455–466.
- Smith JA, Seltzer GO, Farber DL, Rodbell DT, Finkel RC. 2005. Early Local Last Glacial Maximum in the Tropical Andes. *Science* 308: 678–681.
- Soil Survey Staff. 2006. Keys to Soil Taxonomy, 10th edition. *USDA-Natural Resources Conservation Service*, Washington DC (USA).
- Starr MR. 1991. Soil formation and fertility along a 5000 year chronosequence. In: Pulkkinen, E, editor. *Environmental Geochemistry in Northern Europe*. Geological Survey of Finland, Special Paper 9: 99-104.
- Stone JO. 2000. Air pressure and cosmogenic isotope production. *Journal of Geophysical Research* 105 (B10) (23): 753-759.
- Studer M. 2005. Gletschergeschichtliche Untersuchungen und geomorphologische Kartierung im Raum Maloja - Val Forno. Ein Beitrag zur regionalen Landschaftsgeschichte. [Diplomarbeit] Geographisches Institut Universität Zürich.
- Theng BGK, Tate KR, Becker-Heidmann P. 1992. Towards establishing the age, location and identity of the inert organic matter of a spodosol. *Zeitschrift für Pflanzenernährung und Bodenkunde* 155: 181-184.

Table 1 Characteristics of the investigated soils.

Profile	Elevation (m a.s.l.)	Aspect (°N)	Slope (%)	Parent material	Vegetation	WRB (IUSS Working Group, 2006)
S1	2100	60	32	Paragneiss morainic material	<i>Larix decidua</i> / <i>Juniperus communis</i>	Entic Podzol
S2	2230	70	55	Paragneiss morainic material	<i>Rhododendro</i> – <i>vaccinietum extrasilvaticum</i>	Haplic Podzol
S3	2380	320	5	Paragneiss morainic material	<i>Carex curvula</i> / <i>Nardus stricta</i>	Cambic Leptosol
S4	2370	300	10	Paragneiss, inactive rock glacier/solifluction	<i>Carex curvula</i> / <i>Nardus stricta</i>	Brunic Regosol



Table 2 List of samples for surface exposure dating, elevation and latitude of the sample sites, thickness of sample, dip (angle from horizontal) of the surface sample, amount of quartz retrieved from the sample that was used for the measurement of  $^{10}\text{Be}$ , correction factor for topography, production rate,  $^{10}\text{Be}$  measured concentration in the sample, measurement error and  $^{10}\text{Be}$  date.

Sample	Elevation (m asl)	Latitude (°N)	Longitude (°E)	Sample thickness (cm)	Surface dip (°)	Direction of the dip (°)	Shielding correction	Depth corr.	Snow cover* (m)	Local prod. rate (at g <sup>-1</sup> yr <sup>-1</sup> )	$^{10}\text{Be}$ ** (at g <sup>-1</sup> [1E+5])	Error*** (%)	$^{10}\text{Be}$ date (yr)	$^{10}\text{Be}$ cor
B1 (W67)	2247	46.3763	10.7697	3	30	270	0.931	0.975	1.3	27.70	3.23	10.1	11680 ± 1180	13
B2 (W70)	2360	46.3736	10.7883	5	38	270	0.927	0.958	0.7	29.27	3.24	8.5	11110 ± 940	11
B3 (W72)	2456	46.3747	10.7911	5	30	220	0.961	0.958	0.3	32.34	3.15	7.9	9780 ± 770	9
B4 (W73)	2446	46.3750	10.7902	5	16	240	0.982	0.958	0.3	32.84	2.85	7.8	8710 ± 680	8

\* average value of snow cover during 6 months

\*\* using ETH AMS standard S555 ( $^{10}\text{Be}/9\text{Be} = 95.5 \cdot 10^{-12}$  nominal) with a  $^{10}\text{Be}$  half-life of 1.51 Ma

\*\*\* estimated total error including measurement error and the effects of altitude, latitude and topography/depth scaling

Table 3 Physical characteristics of the investigated soils.

Site	Horizons	Depth (cm)	Munsell color (moist)	Skeleton <sup>1)</sup> (%)	Sand (%)	Silt (%)	Clay (%)
S1	AE	0-4	10YR 3/3	5	45.5	28.0	26.5
	BE	4-8	5YR 4/4	11	51.5	28.0	20.5
	Bs1	8-20	7.5YR 4/4	51	57.5	28.6	13.9
	Bs2	20-45	10YR 4/4	45	67.1	27.5	54.0
	BC	45-60	10YR 5/4	34	nd	nd	nd
S2	AE	0-9	7.5YR 2/1	3	83.9	13.6	25.0
	Bhs	9-20	7.5YR 3/3	19	71.7	20.9	74.0
	Bs	20-40	7.5YR 4/3	58	70.9	25.2	39.0
S3	AE1	0-4	10YR 2/3	8	45.7	22.3	32.0
	AE2	4-12	10YR 3/2	21	57.6	21.2	21.2
	Bhs	12-20	10YR 1.7/1	45	63.8	17.2	19.0
S4							

A	0-8	10YR 3/2	0	35.2	49.6	15.2
Bw1	8-20	10YR 4/4	1	40.9	43.7	15.4
Bw2	20-32	10YR 4/4	32	69.2	25.8	50.0
Ab	32-35	10YR 3/3	2	30.9	49.8	19.3
Bb	35-40	10YR 4/4	49	83.9	13.6	25.0

<sup>1)</sup> Size fractions: skeleton = > 2mm; sand = 2000–62µm; silt = 62–2µm; clay = < 2 µm  
nd = not determined

Table 4 Chemical characterisation of the investigated soils.

Site	Horizons	pH (CaCl <sub>2</sub> )	Org. C (%)	Total N (%)	C/N	Al <sub>o</sub> (g/kg)	Fe <sub>o</sub> (g/kg)	Al <sub>d</sub> (g/kg)	Fe <sub>d</sub> (g/kg)
S1	AE	3.7	10.37	0.57	18	1.73	5.57	2.50	15.90
	BE	3.6	6.10	0.29	21	1.91	6.06	2.80	20.50
	Bs1	4.1	3.94	0.18	22	10.27	19.62	14.70	44.10
	Bs2	4.4	1.70	0.07	24	5.84	9.37	7.30	21.40
	BC	4.5	0.75	0.06	12	4.04	1.67	5.60	6.90
S2	AE	3.4	18.46	0.88	21	2.78	5.67	3.94	14.53
	Bhs	3.7	6.38	0.27	24	6.31	24.90	5.96	45.33
	Bs	4.15	2.54	0.12	21	6.41	8.81	10.65	30.13
S3	AE1	3.4	12.49	0.68	18	2.03	2.47	2.80	8.50
	AE2	3.55	4.80	0.22	22	2.48	4.33	3.20	11.00
	Bhs	3.85	7.14	0.31	23	8.30	13.76	14.20	27.10
S4	A	3.85	5.53	0.38	15	3.05	7.05	5.40	21.90
	Bw1	4.00	2.07	0.15	14	2.47	9.61	5.00	30.80
	Bw2	4.1	1.95	0.13	15	1.58	4.21	3.30	20.60
	Ab	3.95	6.20	0.39	16	4.39	6.52	7.50	23.10
	Bbw	4.25	0.91	0.05	18	1.57	3.70	2.70	15.30

Table 5 Calibrated radiocarbon (<sup>14</sup>C) ages (cal BP) of untreated and of H<sub>2</sub>O<sub>2</sub>-treated soil samples.

Site	Soil Type, depth (cm)	Soil horizon	<sup>14</sup> C untreated	<sup>14</sup> C H <sub>2</sub> O <sub>2</sub> (CalPal)	<sup>14</sup> C H <sub>2</sub> O <sub>2</sub> (OxCal)
S1	Entic Podzol				
	0-4	AE	Modern	14683 ± 313	14562 ± 402
	4-8	BE	Modern	17729 ± 237	17310 ± 528
	8-20	Bs1	712 ± 24	11626 ± 213	11623 ± 349
	20-45	Bs2	2929 ± 55	11068 ± 147	11028 ± 242
	45-60	BC	nd	nd	nd
S2	Haplic Podzol				
	0-9	AE	nd	2422 ± 68	2453 ± 246
	9-20	Bhs	nd	nd	nd

S3	20-40	Bs	nd	$11187 \pm 56$	$11105 \pm 280$
	Cambic Leptosol				
	0-4	AE1	nd	$5846 \pm 71$	$5859 \pm 130$
	4-12	AE2	nd	nd	nd
S4	12-20	Bhs	$616 \pm 43$	$10690 \pm 110$	$10754 \pm 319$
	Haplic Regosol				
	0-8	A	nd	$8649 \pm 58$	$8477 \pm 107$
	8-20	Bw1	nd	nd	nd
	20-32	Bw2	nd	$8888 \pm 104$	$8860 \pm 213$
	32-35	Ab	$2594 \pm 101$	$13840 \pm 179$	$13793 \pm 197$
	35-40	Bbw	nd	nd	nd

---

nd = not determined




Figure 1 Location of the investigation site, soil profiles, rock boulders (S = soil profile; B = boulder) and main geomorphologic features.




Figure 2 A) Location of the soil profile S4; B) Soil profile S4 with horizons and buried soil.




Figure 3 Relationship between the obtained ages ( $^{14}\text{C}$  and  $^{10}\text{Be}$ ) and the elevation of the dated objects. Both ages ( $^{14}\text{C}$  and  $^{10}\text{Be}$ ) are referred to 1950 (= 0 BP).




Figure 4 Absolute ages of the investigated sites related to the chronology of the Lateglacial and Holocene glacier and climate variations (according to several authors, e.g. Maisch 1987; Maisch et al. 1999; Kerschner et al. 1999; Ivy-Ochs et al. 2004). All ages ( $^{14}\text{C}$  and  $^{10}\text{Be}$ ) are referred to 1950 (=0 BP).




Figure 5 Location and geomorphic details in the surroundings of the profiles S3, S4 and of the boulders B2, B3 and B4.